arXiv:0902.3626v1 [math.CT] 20 Feb 2009

On pseudocategories in a category with a 2-cell structure

N. Martins-Ferreira

Abstract. For a given (fixed) category, we consider the category of all 2-cell structures (over it) and study some naturality properties. A category with a 2-cell structure is a sesquicategory; we use additive notation for the vertical composition of 2-cells; instead of a law for horizontal composition we consider a relation saying which pairs of 2-cells can be horizontally composed; for a 2-cell structure with every 2-cell invertible, we also consider a notion of commutator, measuring the obstruction for horizontal composition. We compare the concept of naturality in an abstract 2-cell structure with the example of internal natural transformations in a category of the form Cat(**B**), of internal categories in some category **B**, and show that they coincide. We provide a general construction of 2-cell structures over an arbitrary category, under some mild assumptions. In particular, the canonical 2-cell structures over groups and crossed-modules, respectively "conjugations" and "derivations", are instances of these general constructions. We define cartesian 2-cell structure and extend the notion of pseudocategory from the context of a 2-category (as in [**6**]) to the more general context of a sesquicategory. As an example of application we consider pseudocategories in the sesquicategory of abelian chain complexes.

1. Introduction

In this article we use a different notation for the vertical composition of 2-cells: instead of the usual dot '$\cdot$' we use plus '$+$'. To support this we present the following analogy between geometrical vectors in the plane and 2-cells between morphisms

---

2000 *Mathematics Subject Classification.* Primary 18D05, 18D35; Secondary 18E05.

Key words and phrases. sesquicategory, 2-cell structure, cartesian 2-cell structure, natural 2-cell structure, pseudocategory.

This is the result of several talks given by the author in Milan (Oct2006), Coimbra (May2007) and Haute-Bodeux (Jun2007). The author thanks to Professor G. Janelidze for much appreciated help of various kinds.

in a category.



Two geometrical vectors in the plane can be added only if the end point of the second (u as in the picture above) is the starting point of the first one (v as in the picture) and in that case the resulting vector (the sum) goes from the starting point of the second to the end point of the first: exactly the same as with 2-cells

$$\begin{tikzcd}[column sep=large] \cdot \arrow[r, bend left=70, "\mathrm{dom}\,u"{name=A}] \arrow[r, ""{name=B}] \arrow[r, bend right=70, "\mathrm{cod}\,v"'{name=C}] \arrow[Rightarrow, from=A, to=B, "u"] \arrow[Rightarrow, from=B, to=C, "v"] & \cdot & \longmapsto & \cdot \arrow[r, bend left=50, "\mathrm{dom}\,u"{name=D}] \arrow[r, bend right=50, "\mathrm{cod}\,v"'{name=E}] \arrow[Rightarrow, from=D, to=E, "v+u"] & \cdot \, ; \end{tikzcd}$$

In some sense the analogy still holds for scalar multiplication

$$\begin{tikzcd} \cdot \arrow[r, "\lambda"] & \cdot \arrow[r, bend left=50, "\mathrm{dom}\,u"{name=A}] \arrow[r, bend right=50, "\mathrm{cod}\,u"'{name=B}] \arrow[Rightarrow, from=A, to=B, "u"] & \cdot \arrow[r, "\rho"] & \cdot & \longmapsto & \cdot \arrow[r, bend left=50, "\rho\,\mathrm{dom}(u)\lambda"{name=C}] \arrow[r, bend right=50, "\rho\,\mathrm{cod}(u)\lambda"'{name=D}] \arrow[Rightarrow, from=C, to=D, "\rho u\lambda"] & \cdot \, , \end{tikzcd}$$

and for inverses (in the case they exist)

$$\begin{tikzcd} \cdot \arrow[r, bend left=50, "\mathrm{dom}\,u"{name=A}] \arrow[r, bend right=50, "\mathrm{cod}\,u"'{name=B}] \arrow[Rightarrow, from=A, to=B, "u"] & \cdot & \longmapsto & \cdot \arrow[r, bend left=50, "\mathrm{cod}\,u"{name=C}] \arrow[r, bend right=50, "\mathrm{dom}\,u"'{name=D}] \arrow[Rightarrow, from=C, to=D, "-u"] & \cdot \, . \end{tikzcd}$$

Concerning horizontal composition, there is still an analogy with some relevance: it is, in some sense, analogous to the cross product of vectors – in the sense that it raises in dimension (see the introduction of [**1**] and its references for further discussion on this). Given 2-cells, u and v

$$\begin{tikzcd} \cdot \arrow[r, bend left=50, "\mathrm{dom}\,u"{name=A}] \arrow[r, bend right=50, "\mathrm{cod}\,u"'{name=B}] \arrow[Rightarrow, from=A, to=B, "u"] & \cdot \arrow[r, bend left=50, "\mathrm{dom}\,v"{name=C}] \arrow[r, bend right=50, "\mathrm{cod}\,v"'{name=D}] \arrow[Rightarrow, from=C, to=D, "v"] & \cdot \end{tikzcd}$$

the horizontal composition $v \circ u$ should be a 3-cell, from the 2-cell

$$\mathrm{cod}\,(v)\,u + v\,\mathrm{dom}\,(u) \tag{1.1}$$

to the 2-cell

$$v\,\mathrm{cod}\,(u)+\mathrm{dom}\,(v)\,u. \tag{1.2}$$

In some cases (1.1) and (1.2) coincide (as it happens in a 2-category) and this is the reason why one may think of a horizontal composition, but it is an illusion; to overcome this we better consider a relation $v\circ u$ saying that the 2-cell v is natural with respect to u, defined as

$$v\circ u\Longleftrightarrow(1.1)=(1.2)\,,$$

in this sense, the horizontal composition is only defined for those pairs (v,u) that are in relation $v\circ u$, with the composite being then given by either (1.1) or (1.2).

This is a geometrical intuition. An algebraic intuition is also provided in Proposition 1.

This article is organized as follows.

For a fixed category, **C**, we define a 2-cell structure (over **C**, as to make it a sesquicategory) and give a characterization of such a structure as a family of sets, together with maps and actions, satisfying some conditions. It generalizes the characterization of 2-Ab-categories as a family of abelian groups, together with group homomorphisms and laws of composition as given in [**5**] and [**7**] where the (strong) condition

$$D\,(x)\,y=xD\,(y)$$

is no longer required. A useful consequence is that the example of chain complexes, say of order 2, can be considered in this more general setting. Of course, this condition is equivalent to the naturality condition, and the results obtained in [**5**] and [**7**] heavily rest on this assumption, so one must be careful in removing it. For this we introduce and study the concept of a 2-cell being natural with respect to another 2-cell, and the concept of natural 2-cell, as one being natural with respect to all. Next we compare this notions when **C** is a category of the form Cat(**B**), of internal categories in some category **B**, and conclude that if the 2-cell structure is the canonical one (internal transformations, not necessarily natural) then a natural 2-cell corresponds to a natural transformation, and furthermore, it is sufficient to check if a given transformation is natural with respect to a particular 2-cell (from the "category of arrows"), to determine if it is natural.

We give a general process for constructing 2-cell structures in arbitrary categories, and for the purposes of latter discussions we will restrict our study to the 2-cell structures obtained this way. In order to argue that we are not restricting too much, we show that the canonical 2-cell structures over groups and crossed-modules, that are respectively "conjugations" and "derivations", are captured by this construction.

We introduce the notion of cartesian 2-cell structure, in order to consider 2-cells of the form $u\times_w v$ that are used in the coherence conditions involved in a pseudocategory.

At the end we extend the notion of pseudocategory from the context of a 2-category to the more general context of a category with a 2-cell structure (sesquicategory). As an example of application we consider the sesquicategory of abelian chain complexes with homotopies as 2-cells and study pseudocategories in there.

All the notions defined in [**6**]: pseudofunctor, natural and pseudo-natural transformation, modification, may also be extended in this way. However some careful is needed when dealing with coherence issues. For example MacLane's Coherence

Theorem, saying that it suffices to consider the coherence for the *pentagon* and *middle triangle* is no longer true in general, since it uses the fact that α, λ, ρ are natural. One way to overcome this difficulty is to impose the naturality for α, λ, ρ in the definition, so that in [**6**] (introduction, definition of pseudocategory in a 2-category) instead of saying

"...α, λ, ρ are 2-cells (which are isomorphisms)..."

we have to say

"...α, λ, ρ are natural and invertible 2-cells ..."

We will not study deeply all the consequences of this. Instead we will restrict ourselves to the study of 2-cell structures such that all 2-cells are invertible (since the main examples are groups, abelian groups, 1-chain complexes and crossed modules) and hence the question of α, λ, ρ being invertible becomes intrinsic to the 2-cell structure. The issue of naturality is more delicate. To prove the results in [**5**], [**7**] and [**9**], we will only need λ and ρ to be natural with respect to each other, that is

$$\lambda \circ \lambda, \lambda \circ \rho, \rho \circ \lambda, \rho \circ \rho.$$

If interested in the Coherence Theorem, we can always use the reflection

$$\text{2-cellstruct}(\mathbf{C}) \xrightarrow{I} \text{nat-2-cellstruct}(\mathbf{C})$$

of the category of 2-cell structures over $\mathbf{C}$ (sesquicategories "with base $\mathbf{C}$"), into the subcategory of natural 2-cell structures over $\mathbf{C}$ (2-categories "with base $\mathbf{C}$"), sending each 2-cell structure to its "naturalization"; which, if $\mathbf{C} = \mathbf{1}$, becomes the familiar reflection of monoids into commutative monoids

$$\text{Mon} \xrightarrow{I} \text{CommMon}$$

and if restricting further to invertible 2-cells gives the reflection

$$\text{Grp} \xrightarrow{I} \text{Ab}$$

of groups into abelian groups.

All these considerations are to be developed in some future work. This paper is the starting point for a systematic study of internal categorical structures in a category with a given 2-cell structure, and also to investigate how these categorical structures are changed when the given 2-cell structure over the (fixed) base category also changes. For example, a pseudo category, in a category with the discrete 2-cell structure is an internal category, while if the 2-cell structure is the codiscrete one, it becomes simply a precategory.

2. 2-cell structures and sesquicategories

Let $\mathbf{C}$ be a fixed category.

Definition 1 (2-cell structure). *A 2-cell structure over* $\mathbf{C}$ *is a system*

$$\mathbf{H} = (H, \text{dom}, \text{cod}, 0, +)$$

where

$$H : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow Set$$

is a functor and

$$\begin{tikzcd} H \times_{\hom} H \arrow[r, "+"] & H \arrow[r, shift left=3, "\mathrm{dom}"] \arrow[r, shift right=3, "\mathrm{cod}"'] & \hom_{\mathbf{C}} \arrow[l, "0"'] \end{tikzcd}$$

are natural transformations, such that

$$(\hom_{\mathbf{C}}, H, \mathrm{dom}, \mathrm{cod}, 0, +)$$

is a category object in the functor category $Set^{C^{op} \times C}$ *or, in other words, an object in* $Cat\left(Set^{C^{op} \times C}\right)$.

Proposition 1. *Giving a 2-cell structure over a category* $\mathbf{C}$*, is to give, for every pair* (A, B) *of objects in* $\mathbf{C}$*, a set* $H\,(A, B)$*, together with maps*

$$\begin{tikzcd} H\,(A,B) \times_{\hom(A,b)} H\,(A,B) \arrow[r, "+"] & H\,(A,B) \arrow[r, shift left=3, "\mathrm{dom}"] \arrow[r, shift right=3, "\mathrm{cod}"'] & \hom\,(A,B)\,, \arrow[l, "0"'] \end{tikzcd}$$

and actions

$$\begin{array}{ccc} H\,(B,C) \times \hom\,(A,B) & \longrightarrow & H\,(A,C) \\ (x,f) & \longmapsto & xf \\ \hom\,(B,C) \times H\,(A,B) & \longrightarrow & H\,(A,C) \\ (g,y) & \longmapsto & gy \end{array}$$

satisfying the following conditions

$$\begin{gathered} \mathrm{dom}\,(gy) = g\,\mathrm{dom}\,(y)\ \ ,\ \mathrm{dom}\,(xf) = \mathrm{dom}\,(x)\,f \\ \mathrm{cod}\,(gy) = g\,\mathrm{cod}\,(y)\ \ ,\ \mathrm{cod}\,(xf) = \mathrm{cod}\,(x)\,f \\ g0_f = 0_{gf} = 0_g f \\ (x + x')\,f = xf + x'f\ ,\ g\,(y + y') = gy + gy' \end{gathered} \tag{2.1}$$

$$\begin{aligned} g'\,(gy) &= (g'g)\,y\ ,\ (xf)\,f' = x\,(ff') \\ g'\,(xf) &= (g'x)\,f \\ 1_C x &= x = x1_B \end{aligned} \tag{2.2}$$

$$\begin{aligned} \mathrm{dom}\,(0_f) &= f = \mathrm{cod}\,(0_f) \\ \mathrm{dom}\,(x + x') &= x'\ ,\ \mathrm{cod}\,(x + x') = x \\ 0_{\mathrm{cod}\,x} + x &= x = x + 0_{\mathrm{dom}\,x} \\ x + (x' + x'') &= (x + x') + x''. \end{aligned} \tag{2.3}$$

Proof. For every $f : A' \longrightarrow A$, $g : B \longrightarrow B'$ and $x \in H\,(A, B)$, write

$$H\,(f, g)\,(x) = gxf$$

and it is clear that the set of conditions (2.1) asserts the naturality of $\mathrm{dom}, \mathrm{cod}, 0, +$; the set of conditions (2.2) asserts the functoriality of H and the set of conditions (2.3) asserts the axioms for a category. □

Definition 2 (sesquicategory). *A* sesquicategory *is a pair* $(\mathbf{C}, \mathbf{H})$ *where* $\mathbf{C}$ *is a category and* $\mathbf{H}$ *a 2-cell structure over it.*

Observation: A sesquicategory, as defined, is the same as a sesquicategory in the sense of Ross Street ([**10**],[**11**],[**12**]), that is, a category $\mathbf{C}$ together with a functor $\mathbf{H}$ into Cat, such that the restriction to Set gives $\hom_{\mathbf{C}}$, as displayed in the following picture

$$\begin{tikzcd}
 & Cat\ . \arrow[d] \\
\mathbf{C}^{op}\times\mathbf{C} \arrow[ur, "\mathbf{H}"] \arrow[r, "\hom"'] & Set
\end{tikzcd}$$

Proposition 2. *A category* $\mathbf{C}$ *with a 2-cell structure*

$$\mathbf{H} = (H, \mathrm{dom}, \mathrm{cod}, 0, +)\,,$$

is a 2-category if and only if the naturality condition

$$\text{(naturality condition)} \qquad \mathrm{cod}\,(x)\,y + x\,\mathrm{dom}\,(y) = x\,\mathrm{cod}\,(y) + \mathrm{dom}\,(x)\,y$$

holds for every $x \in H\,(B,C)\,,\ y \in H\,(A,B)\,,$ *and every triple of objects* (A,B,C) *in* $\mathbf{C}$*, as displayed in the diagram below*

$$\begin{tikzcd}
A \arrow[r, bend left=50, "\mathrm{dom}\, y"] \arrow[r, bend right=50, "\mathrm{cod}\, y"'] \arrow[r, phantom, "\Downarrow y"] & B \arrow[r, bend left=50, "\mathrm{dom}\, x"] \arrow[r, bend right=50, "\mathrm{cod}\, x"'] \arrow[r, phantom, "\Downarrow x"] & C\ .
\end{tikzcd}$$

Proof. If $\mathbf{C}$ is a 2-category, the naturality condition follows from the horizontal composition of 2-cells and, conversely, given a 2-cell structure over $\mathbf{C}$, in order to make it a 2-category one has to define a horizontal composition and it is defined as

$$x \circ y = \mathrm{cod}\,(x)\,y + x\,\mathrm{dom}\,(y)$$

or

$$x \circ y = x\,\mathrm{cod}\,(y) + \mathrm{dom}\,(x)\,y$$

provided the naturality condition is satisfied for every appropriate x, y. The middle interchange law also follows from the naturality condition. □

It may happen that the naturality condition does not hold for all possible x and y, but only for a few; thus the following definitions.

Let $\mathbf{C}$ be a category and $(H, \mathrm{dom}, \mathrm{cod}, 0, +)$ a 2-cell structure over it.

Definition 3. *A 2-cell* $\delta \in H\,(A,B)$ *is* natural with respect to *a 2-cell* $z \in H\,(X,A)$*, when*

$$\mathrm{cod}\,(\delta)\,z + \delta\,\mathrm{dom}\,(z) = \delta\,\mathrm{cod}\,(z) + \mathrm{dom}\,(\delta)\,z;$$

in that case one writes $\delta \circ z$*.*

Definition 4. *A 2-cell* $\delta \in H\,(A,B)$ *is* natural *when it is natural with respect to all possible* $z \in H\,(X,A)$ *for all* $X \in \mathbf{C}$*, i.e.,* δ *is a natural 2-cell if and only if* $\delta \circ z$ *for all possible* z*.*

3. Examples of application

We shall now see how the above notions of naturality are related, in the case where $\mathbf{C} = Cat\,(\mathbf{B})$ for some category $\mathbf{B}$, with the 2-cell structure given by the internal (natural) transformations.

3.1. The canonical 2-cell structure of Cat(B). Consider $\mathbf{C} = Cat\left(\mathbf{B}\right)$ the category of internal categories in some category $\mathbf{B}$. The objects are[1] (see also [**4**], p.267)

$$A = \left(A_0, A_1, d, c, e, m\right),\ B = \left(B_0, B_1, d, c, e, m\right),\ ...$$

and morphisms

$$f = \left(f_1, f_0\right) : A \longrightarrow B, ...$$

In this case, we have a canonical 2-cell structure, given by the internal transformations (not necessarily natural) and it is as follows:

$$\begin{aligned} H\left(A,B\right) &= \left\{\left(k,t,h\right) \mid t : A_0 \longrightarrow B_1;\ h,k \in \hom_{\mathbf{C}}\left(A,B\right);\ dt = h_0, ct = k_0\right\} \\ H\left(f,g\right)\left(k,t,h\right) &= \left(gkf, g_1tf_0, ghf\right) \\ \operatorname{dom}\left(k,t,h\right) &= h \\ \operatorname{cod}\left(k,t,h\right) &= k \\ 0_h &= \left(h, eh_0, h\right) \\ \left(k,t,h\right) + \left(h,s,l\right) &= \left(k, m\left\langle t,s\right\rangle, l\right) \end{aligned}$$

where $f : A' \longrightarrow A, g : B \longrightarrow B', h,k,l : A \longrightarrow B$ are morphisms in $Cat\left(\mathbf{B}\right)$ and $t,s : A_0 \longrightarrow B_1$ are morphisms in $\mathbf{B}$.

Observe that, in particular, for every $A = \left(A_0, A_1, d, c, e, m\right)$ there is $A^{\rightarrow} = \left(A_1, A_1, 1, 1, 1, 1\right)$ and the two morphisms

$$d^{\rightarrow} = \left(ed, d\right) : A^{\rightarrow} \longrightarrow A$$

and

$$c^{\rightarrow} = \left(ec, c\right) : A^{\rightarrow} \longrightarrow A.$$

Proposition 3. *In the context of the above setting, a 2-cell* $\mathbf{t} = \left(k,t,h\right) \in H\left(A,B\right)$ *is an internal natural transformation* $\mathbf{t} : h \longrightarrow k$ *if and only if it is natural with respect to the 2-cell*

$$\left(c^{\rightarrow}, 1_{A_1}, d^{\rightarrow}\right) \in H\left(A^{\rightarrow}, A\right).$$

Proof. Consider $\mathbf{t} = \left(k,t,h\right) \in H\left(A,B\right)$ and $\mathbf{z} = \left(g,z,f\right) \in H\left(X,A\right)$,

$$\begin{tikzcd}
\cdots & X_1 \arrow[r, shift left=2] \arrow[r, shift right=2] \arrow[d, shift right, "f_1"'] \arrow[d, shift left, "g_1"] & X_0 \arrow[l] \arrow[dl, "z"'] \arrow[d, shift right, "f_0"'] \arrow[d, shift left, "g_0"] \\
\cdots & A_1 \arrow[r, shift left=2] \arrow[r, shift right=2] \arrow[d, shift right, "h_1"'] \arrow[d, shift left, "k_1"] & A_0 \arrow[l] \arrow[dl, "t"'] \arrow[d, shift right, "h_0"'] \arrow[d, shift left, "k_0"] \\
\cdots & B_1 \arrow[r, shift left=2] \arrow[r, shift right=2] & B_0 \arrow[l]
\end{tikzcd}$$

by definition

$$\begin{aligned} \mathbf{t} \circ \mathbf{z} &\Leftrightarrow \left(kg, k_1z, kf\right) + \left(kf, tf_0, hf\right) = \left(kg, tg_0, hg\right) + \left(hg, h_1z, hf\right) \\ &\Leftrightarrow \left(kg, m\left\langle k_1z, tf_0\right\rangle, hf\right) = \left(kg, m\left\langle tg_0, h_1z\right\rangle, hf\right) \\ &\Leftrightarrow m\left\langle k_1z, tf_0\right\rangle = m\left\langle tg_0, h_1z\right\rangle \qquad (3.1) \end{aligned}$$

and also by definition t is an internal natural transformation when

$$m\left\langle k_1, td\right\rangle = m\left\langle tc, h_1\right\rangle \qquad (3.2)$$

---

[1] An internal category is a system $\left(C_0, C_1, d, c, e, m\right)$ where $d, c : C_1 \longrightarrow C_0, e : C_0 \longrightarrow C_1, m : C_1 \times_{C_0} C_1 \longrightarrow C_1$ satisfying $de = 1_{C_0} = ec$, plus the usual axioms for preservation of domain, d, codomain, c, identity (e) for composition (m) and associativity.

which is equivalent to saying that (k, t, h) is natural relative to $(c^{\rightarrow}, 1_{A_1}, d^{\rightarrow})$, as displayed below

$$\begin{tikzcd}
\cdots & A_1 \arrow[r, equal] \arrow[d, shift right, "ed"'] \arrow[d, shift left, "ec"] & A_1 \arrow[dl, "1"'] \arrow[d, shift right, "d"'] \arrow[d, shift left, "c"] & . \\
\cdots & A_1 \arrow[r, shift left=2] \arrow[r, shift right=2] & A_0 \arrow[l]
\end{tikzcd}$$

$\square$

Corollary 1. *Every internal natural transformation is a natural 2-cell.*

Proof. Simply observe that

$$(3.2) \Longrightarrow (3.1)$$

since

$$\begin{aligned}
m\langle k_1, td\rangle z &= m\langle tc, h_1\rangle z \\
m\langle k_1 z, tdz\rangle &= m\langle tcz, h_1 z\rangle \\
m\langle k_1 z, tf_0\rangle &= m\langle tg_0, h_1 z\rangle .
\end{aligned}$$

$\square$

The notion of a category with a 2-cell structure, besides giving a simple characterization of a 2-category as

"2-category" = "sesquicategory" + "naturality condition";

it also provides a powerful tool to construct examples in arbitrary situations.

3.2. Abstract 2-cells, and conjugations.

Consider $\mathbf{C}$ a category and

$$H : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow Mon$$

a functor into Mon, the category of monoids, together with a natural transformation

$$D : UH \times \hom_{\mathbf{C}} \longrightarrow \hom_{\mathbf{C}}$$

(where $U : Mon \longrightarrow Set$ denotes the forgetful functor) satisfying

$$\begin{aligned}
D(0, f) &= f \\
D(x' + x, f) &= D(x', D(x, f))
\end{aligned}$$

for all $f : A \longrightarrow B$ in $\mathbf{C}$ and $x', x \in H(A, B)$, with 0 the zero of the monoid $H(A, B)$ considered in additive notation.

A 2-cell structure in $\mathbf{C}$ is now given as

$$\begin{tikzcd}
A \arrow[r, bend left=50, "f"{name=U}] \arrow[r, bend right=50, "D(x,f)"'{name=L}] & B
\arrow[Rightarrow, from=U, to=L, "(x,f)" description]
\end{tikzcd}$$

with vertical composition

$$\begin{tikzcd}
A \arrow[rr, bend left=70, "f"{name=U}] \arrow[rr, "D(x,f)"{name=M, description}] \arrow[rr, bend right=70, "{D(x',D(x,f))}"'{name=L}] & & B
\arrow[Rightarrow, from=U, to=M, "{(x,f)}"] \arrow[Rightarrow, from=M, to=L, "{(x',D(x,f))}"]
\end{tikzcd}$$

$$(x', D(x, f)) + (x, f) = (x' + x, f)$$

(well defined because $D(x' + x, f) = D(x', D(x, f))$), with identity 2-cells

$$\begin{tikzcd}
A \arrow[r, bend left=50, "f"{name=U}] \arrow[r, bend right=50, "f"'{name=L}] & B
\arrow[Rightarrow, from=U, to=L, "{(0,f)}"]
\end{tikzcd}$$

well defined because $D(0, f) = f$, and the left and right actions of morphisms in 2-cells,

$$\begin{tikzcd}
A' \arrow[r, "h"] & A \arrow[r, bend left=50, "f"{name=U}] \arrow[r, bend right=50, "{D(x,f)}"'{name=L}] & B \arrow[r, "g"] & B'
\arrow[Rightarrow, from=U, to=L, "{(x,f)}"]
\end{tikzcd}$$

$$g(x, f)h = (gxf, gfh) = (H(h, g)(x), gfh).$$

If in addition,

$$D(y, g)x + yf = yD(x, f) + gx \tag{3.3}$$

for all x, y, f, g pictured as

$$\begin{tikzcd}
A \arrow[r, bend left=50, "f"{name=U1}] \arrow[r, bend right=50, "{D(x,f)}"'{name=L1}] & B \arrow[r, bend left=50, "g"{name=U2}] \arrow[r, bend right=50, "{D(y,g)}"'{name=L2}] & C
\arrow[Rightarrow, from=U1, to=L1, "{(x,f)}"] \arrow[Rightarrow, from=U2, to=L2, "{(y,g)}"]
\end{tikzcd},$$

then the result is a 2-category.

3.2.1. *The case of Groups.* In the case of $\mathbf{C} = Grp$ the category of groups and group homomorphisms, the construction above is so general that it includes the canonical 2-cells that are obtained if considering each group as a one object groupoid and each group homomorphism as a functor. In that case, as it is well known, a 2-cell

$$t : f \longrightarrow g$$

from the homomorphism f to the homomorphism g, both from the group A to the group B, is an element $t \in B$ such that

$$tf(x) = g(x)t \quad , \quad \text{for all } x \in A.$$

Now, given t and f, the homomorphism g is uniquely determined as

$$g\left(x\right)=tf\left(x\right)t^{-1}={}^{t}f\left(x\right),$$

and hence, this particular 2-cell structure over *Grp* is an instance of Example 3.2 with *Grp* instead of Mon.

To see this just consider H the functor that projects the second argument

$$\begin{gathered} H:Grp^{op}\times Grp\longrightarrow Grp \\ \left(A,B\right)\longmapsto B \end{gathered}$$

and

$$\begin{gathered} D:B\times\hom\left(A,B\right)\longrightarrow\hom\left(A,B\right) \\ \left(t,f\right)\longmapsto {}^{t}f \end{gathered}$$

and it is a straightforward calculation to check that

$$\begin{array}{rcl} D\left(0,f\right) & = & f \\ D\left(t+t',f\right) & = & D\left(t,D\left(t',f\right)\right) \end{array}$$

and also, since condition (3.3) is satisfied, the 2-cell structure is natural.

In some cases, the above construction may even be pushed further.

3.3. Abstract 2-cells, and derivations. Suppose the functor

$$\hom_{\mathbf{C}}:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Set$$

may be extended to Mon, that is, there is a functor (denote it by *map*, and think of the underlying map of a homomorphism)

$$map:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Mon\xrightarrow{U}Set$$

with $\hom\subseteq Umap$, in the sense that $\hom\left(A,B\right)\subseteq Umap\left(A,B\right)$ naturally for every $A,B\in\mathbf{C}$;
Now, given any functor

$$K:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Mon$$

and any natural transformation

$$D:K\longrightarrow map,$$

define

$$\begin{array}{rcl} H\left(A,B\right) & = & \left\{\left(x,f\right)\in UK\left(A,B\right)\times\hom\left(A,B\right)\mid D\left(x\right)+f\in\hom\left(A,B\right)\right\} \\ H\left(h,g\right)\left(x,f\right) & = & \left(gxh,gfh\right) \end{array}$$

and obtain a functor $H:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Set$. With obvious $\mathrm{dom},\mathrm{cod},0,+$, a 2-cell structure in $\mathbf{C}$ is obtained as follows

$$\begin{tikzcd} A \arrow[r, bend left=50, "f"{name=U}] \arrow[r, bend right=50, "D(x)+f"'{name=D}] & B \arrow[Rightarrow, from=U, to=D, "{(x,f)}"] \end{tikzcd}$$

where $\left(x,f\right)\in H\left(A,B\right)$,
vertical composition: $\left(x',D\left(x\right)+f\right)+\left(x,f\right)=\left(x'+x,f\right)$

identity: $(0,f)$
left and right actions: $g\,(x,f)\,h=(gxh,gfh)$.
If in addition the property

$$D\,(y)\,x+gx+yf=yD\,(x)+yf+gx \tag{3.4}$$

is satisfied for all $(x,f)\in H_1\,(A,B)$ and $(y,g)\in H_1\,(A,C)$, then the resulting structure is a 2-category.

3.3.1. *The case of crossed modules.* In the case $\mathbf{C}$=X-Mod, the category of crossed modules, we have the canonical 2-cell structure given by derivations, and it is an instance of the above construction with Grp instead of Mon:

The objects in X-Mod are of the form

$$A=\left(X\xrightarrow{d}B,\varphi:B\longrightarrow Aut\,(X)\right)$$

where $d:X\longrightarrow B$ is a group homomorphism, together with a group action of B in X denoted by $b\cdot x$ satisfying

$$\begin{aligned} d\,(b\cdot x) &= bd\,(x)\,b^{-1}\\ d\,(x)\cdot x' &= x+x'-x; \end{aligned}$$

a morphism $f:A\longrightarrow A'$ in X-Mod is of the form

$$f=(f_1,f_0)$$

where $f_1:X\longrightarrow X'$ and $f_0:B\longrightarrow B'$ are group homomorphisms such that

$$f_0d=d'f_1$$

and

$$f_1\,(b\cdot x)=f_0\,(b)\cdot f_1\,(x)\,.$$

Clearly there are functors

$$map:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Grp$$

sending (A,A') to the group of pairs (f_1,f_0) of maps (not necessarily homomorphisms) $f_1:UX\longrightarrow UX'$ and $f_0:UB\longrightarrow UB'$ such that

$$f_0d=d'f_1,$$

with the group operation defined componentwise

$$(f_1,f_0)+(g_1,g_0)=(f_1+g_1,f_0+g_0)\,.$$

Also there is a functor

$$M:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Grp$$

sending (A,A') to the group $M\,(A,A')=\{t\mid t:UB\longrightarrow UX'\text{ is a map}\}$, and a natural transformation

$$D:M\longrightarrow map$$

defined by

$$D\,(A,A')\,(t)=(td,dt)\,.$$

Now, define $H\,(A,A')$ as

$$\{(t,f)\mid t\in M\,(A,A')\ ,\ f=(f_1,f_0):A\longrightarrow A'\ ,\ (td+f_1,dt+f_0)\in\hom\,(A,A')\}\,.$$

It is well known that the map $t:B\longrightarrow X'$ is such that

$$t\,(bb')=t\,(b)+f_0\,(b)\cdot t\,(b')\quad,\quad\text{for all }b,b'\in B,$$

while $(td+f_1, dt+f_0) \in \hom(A, A')$ asserts that the pair $(td+f_1, dt+f_0)$ is a morphism of crossed modules

$$\begin{tikzcd}
X \arrow[r, "d"] \arrow[d, "td+f_1"'] & B \arrow[d, "dt+f_0"] \\
X' \arrow[r, "d"] & B'
\end{tikzcd} \tag{3.5}$$

and it is equivalent to

- $dt+f_0$ is a homomorphism of groups
$$dt(bb') = d\left(t(b) + f_0(b) \cdot t(b')\right)$$
- $td+f_1$ is a homomorphism of groups
$$t\left(d(x)\, d(x')\right) = t(dx) + f_0 d(x) \cdot td(x')$$
- the square (3.5) commutes, which is trivial because $(f_1, f_0) \in \hom(A, A')$
- $(td+f_1)$ preserves the action of $(dt+f_0)$
$$t\left(bd(x)\, b^{-1}\right) = t(b) + f_0(b) \cdot t(d(x)) + f_0\left(bd(x)\, b^{-1}\right) \cdot (-t(b)).$$

3.4. Abstract 2-cells and homotopies. In particular, if $\mathbf{C}$ is an Ab-category, a 2-Ab-category as defined in [**5**] and [**7**] is obtained in this way; in that case the functor hom is in fact a functor

$$\hom : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow Ab.$$

Giving a 2-cell structure is then to give a functor (usually required to be an Ab-functor) $H : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow Ab$, and a natural transformation $D : H \longrightarrow \hom$. This 2-cell structure makes $\mathbf{C}$ a 2-category (in fact a 2-Ab-category) if in addition the condition (3.4) is satisfied, which in the abelian context simplifies to $D(y)\,x = yD(x)$. Furthermore, as proved in [**5**], every 2-cell structure (if enriched in Ab) is obtained in this way.

Of course, these considerations are also valid for any monoidal category $\mathbf{V}$, except that in general not all 2-cell structures are obtained in this way.

3.4.1. *The case of Abelian Chain Complexes.* The example of abelian chain complexes (say of order 2 to simplify notation) is self explanatory (see also [**13**],[**14**] and references there). We have objects, morphisms and 2-cells (homotopies) as displayed

$$\begin{tikzcd}
A_2 \arrow[r, "d"] \arrow[d, shift right, "f_2"'] \arrow[d, shift left, "g_2"] & A_1 \arrow[r, "d"] \arrow[d, shift right, "f_1"'] \arrow[d, shift left, "g_1"] \arrow[dl, "t_2"'] & A_0 \arrow[d, shift right, "f_0"'] \arrow[d, shift left, "g_0"] \arrow[dl, "t_1"'] \\
A'_2 \arrow[r, "d"] & A'_1 \arrow[r, "d"] & A'_0
\end{tikzcd} \tag{3.6}$$

with

$$\begin{aligned}
dt_1 &= g_0 - f_0 \\
t_1 d + dt_2 &= g_1 - f_1 \\
t_2 d &= g_2 - f_2
\end{aligned}$$

or equivalently

$$\begin{aligned} g_0 &= dt_1 + f_0 \\ g_1 &= t_1 d + dt_2 + f_1 \\ g_2 &= t_2 d + f_2 \end{aligned}$$

and hence we have, for $\mathbf{C}$ =2-Ch(Ab), the functor

$$H : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow Ab$$

sending the pair of objects (A, A') to the abelian group of pairs (t_2, t_1); and the natural transformation

$$D : H \longrightarrow \text{hom}$$

sending a pair (t_2, t_1) as above to the triple $(t_2 d, t_1 d + dt_2, dt_1)$ displayed as follows

$$\begin{tikzcd} A_2 \arrow[r, "d"] \arrow[d, "t_2 d"'] & A_1 \arrow[r, "d"] \arrow[d, "t_1 d + dt_2"'] & A_2 \arrow[d, "dt_1"'] \\ A'_2 \arrow[r, "d"] & A'_1 \arrow[r, "d"] & A'_0 \end{tikzcd} .$$

This is in fact an instance of the above construction, however, condition

$$D(x)y = xD(y)$$

is not satisfied in general, and it becomes, for $x = (t_2, t_1)$, $y = (s_2, s_1)$

$$(t_2 d s_2, dt_2 s_1 + t_1 d s_1) = (t_2 d s_2 + t_2 s_1 d, t_1 d s_1)$$

which holds if $t_2 s_1 = 0$, but not in general.

The commutator (see below) in this case is

$$[x, y] = (-t_2 s_1 d, dt_2 s_1)\,.$$

3.5. Abstract 2-cells and commutators. Previous examples apply to arbitrary (even large) categories, provided they admit the functors and the natural transformations as specified. Interesting examples also appear if one tries to particularize the category $\mathbf{C}$. For example if $\mathbf{C}$ has only one object, or if it is a preorder; the first case gives something that particularizes to a (strict) monoidal category (with fixed set of objects) in the presence of the naturality condition; while the second case gives something that particularizes to an enriched category over monoids. The simplest case, when $\mathbf{C}=\mathbf{1}$, gives Monoids and Commutative Monoids under the naturality condition; so in particular, if considering only invertible 2-cell structures the result is Groups and Abelian Groups, respectively.

The well known reflection

$$Gr \xrightarrow{I} Ab,$$

accordingly to G. Janelidze, generalizes to a reflection

$$\text{2-cellstruct}(\mathbf{C}) \xrightarrow{I} \text{nat-2-cellstruct}(\mathbf{C})$$

from the category of 2-cell structures over $\mathbf{C}$, into the subcategory of natural 2-cell structures over $\mathbf{C}$, sending each 2-cell structure to its "naturalization"; and, under the assumption that all the 2-cells are invertible, one may consider for each

$$\begin{tikzcd} A \arrow[r, bend left=50, ""{name=U1, below}] \arrow[r, bend right=50, ""{name=D1, above}] & B \arrow[r, bend left=50, ""{name=U2, below}] \arrow[r, bend right=50, ""{name=D2, above}] & C \arrow[Rightarrow, from=U1, to=D1, "y"] \arrow[Rightarrow, from=U2, to=D2, "x"] \end{tikzcd}$$

the commutator

$$\begin{aligned} [x,y] &= (c_1 + d_2 - d_1 - c_2)(x,y) \\ &= c_1(x,y) + d_2(x,y) - d_1(x,y) - c_2(x,y) \end{aligned}$$

where

$$\begin{aligned} c_1(x,y) &= \operatorname{cod}(x)\,y\ ,\ c_2(x,y) = x\operatorname{cod}(y) \\ d_1(x,y) &= \operatorname{dom}(x)\,y\ ,\ d_2(x,y) = x\operatorname{dom}(y)\,, \end{aligned}$$

and the comparison with $0_{\operatorname{cod}(x)\operatorname{cod}(y)}$ tell us the obstruction that x and y offer to be horizontally composed.

We will not developed this concept further, at the moment we are only observing that in the case of $\mathbf{C}$ being an Ab-category (see [**5**],[**7**] and Section 3.4 above) the notion of commutator reduces to

$$[x,y] = D(x)\,y - xD(y)\,.$$

In fact the notion of 2-Ab-category (as introduced in [**5**]) may be pushed further in the direction of a sesquicategory enriched in any category $\mathbf{A}$ with a "forgetful" functor into Sets.

It is a simple generalization of Section 3.3 above and it is as follows.

For a category $\mathbf{A}$ with a "forgetful" functor into Sets, $U : \mathbf{A} \longrightarrow$Sets, assume the existence of a functor

$$map : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow \mathbf{A}$$

such that

$$\hom_C(A,B) \subseteq Umap(A,B)$$

(as in Section 3.3).

If $\mathbf{A}$ were monoidal and $\mathbf{C}$ a category enriched in $\mathbf{A}$ then we would always be in the above conditions, simply by choosing $map = \hom$. It is then reasonably to say that, in this more general context, the category $\mathbf{C}$ is weakly enriched in $\mathbf{A}$ (for example, in this sense, Groups are weakly enriched in Groups, and every algebraic structure is weakly enriched in itself). In this conditions, we may be interested in considering only 2-cell structures over $\mathbf{C}$ that are "weakly enriched" in $\mathbf{A}$ in the same way as $\mathbf{C}$ is. This concept is obtained if considering only the 2-cell structures that are given by

$$H(A,B) = \{x \in UM(A,B) \mid U\operatorname{dom} x, U\operatorname{cod} x \in \hom(A,B)\}$$

for some $M, \operatorname{dom}, \operatorname{cod}$ being part of an internal category object in $\mathbf{A}^{\mathbf{C}^{op}\times\mathbf{C}}$, of the form

$$\begin{tikzcd} M \times_{map} M \arrow[r, "+"] & M \arrow[r, shift left=2, "\operatorname{dom}"] \arrow[r, shift right=2, "\operatorname{cod}"'] & map \arrow[l, "0"' description] \end{tikzcd},$$

with the obvious restrictions after applying U.

It is interesting now to observe that in the case of $\mathbf{A} = Grp$ the result of this is precisely the construction of Section 3.3. If $\mathbf{A}$ =Ab and also requiring M to be an Ab-functor, then the result is a 2-Ab-category if also adding the condition

$$D(x)\,y = xD(y)$$

for all appropriate x and y.

3.5.1. *The one-object case.* In the case of a one object category, we may identify it with a monoid, say M and hence giving a 2-cell structure (let us say in the context of Section 3.2 with Ab instead of Mon, for simplicity) over it is to give (see also Proposition 1) an abelian group H (where M acts on the left and on the right) and a map

$$D : H \times M \longrightarrow M$$

satisfying

$$\begin{aligned} D(0, f) &= f \\ D(x' + x, f) &= D(x', D(x, f)) \end{aligned}$$

which is simply an action of the group H on the monoid M. Note that by naturality of D we also have $gD(x, f)\,h = D(gxf, gfh)$. The commutator in this case is given by

$$[x, y] = D(y, g)\,x + yf - yD(x, f) - gx.$$

In the case D is the trivial action we obtain the familiar notion of M semimodule.

4. Morphisms between 2-cell structures and cartesian 2-cell structures

For a fixed category $\mathbf{C}$, there is the category 2-cellstruct($\mathbf{C}$) of all possible 2-cell structures over $\mathbf{C}$, as well as the subcategory nat-2-cellstruct($\mathbf{C}$) of natural 2-cell structures over C and inv-2-cellstruct(C) of all the invertible 2-cell structures over C. The category 2-cellstruct($\mathbf{C}$) has a initial object (the discrete 2-cell structure) and a terminal object (the codiscrete 2-cell structure). If C is of the form Cat(B) for some category B, it also has the canonical 2-cell structure of internal transformations and the canonical natural 2-cell structure of internal natural transformations.

4.1. The category of 2-cell structures over a fixed category C.

The objects of 2-cellstruct($\mathbf{C}$) are of the form

$$\mathbf{H} = (H, \mathrm{dom}, \mathrm{cod}, 0, +)$$

where

$$H : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow Set$$

is a functor and

$$\begin{tikzcd} H \times_{\mathrm{hom}} H \arrow[r, "+"] & H \arrow[r, shift left=3, "\mathrm{dom}"] \arrow[r, shift right=3, "\mathrm{cod}"'] & \mathrm{hom}_{\mathbf{C}} \arrow[l, "0"'] \end{tikzcd}$$

are natural transformations, such that

$$(\mathrm{hom}_{\mathbf{C}}, H, \mathrm{dom}, \mathrm{cod}, 0, +)$$

is a category object in the functor category $\mathrm{Set}^{C^{op}\times C}$,or , in other words, is an object in $\mathrm{Cat}\big(Set^{C^{op}\times C}\big)$.
A morphism $\varphi:\mathbf{H}\longrightarrow\mathbf{H}'$ is a natural transformation

$$\varphi:H\longrightarrow H'$$

such that

$$\begin{aligned}\mathrm{dom}'\,\varphi &= \mathrm{dom}\\ \mathrm{cod}'\,\varphi &= \mathrm{cod}\\ \varphi 0 &= 0'\\ \varphi + &= +'\,(\varphi\times\varphi)\,.\end{aligned}$$

We will often write simply H to refer to a 2-cell structure, whenever confusion is unlikely to appear.

The purpose of describing 2-cellstruct($\mathbf{C}$), the category of all 2-cell structures over a given category $\mathbf{C}$, is the study of pseudocategories in $\mathbf{C}$. The notion of pseudocategory in a category $\mathbf{C}$ depends of the 2-cell structure considered over $\mathbf{C}$. For example, a pseudocategory in $\mathbf{C}$ with the codiscrete 2-cell structure is a precategory, while, if considering the discrete 2-cell structure, it is an internal category. It seems to be interesting to study, for a given category $\mathbf{C}$, how the notion of pseudocategory changes from a precategory to an internal category by changing the 2-cell structure considered over $\mathbf{C}$. This topic was studied in [**9**] for the case of weakly Mal'cev sesquicategories.

Also, every morphism

$$\varphi:H\longrightarrow H' \tag{4.1}$$

in 2-cellstruct($\mathbf{C}$) induces a functor

$$PsCat\,(\mathbf{C},H)\longrightarrow PsCat\,(\mathbf{C},H') \tag{4.2}$$

from pseudocategories in $\mathbf{C}$ relative to the 2-cell structure H to pseudocategories in $\mathbf{C}$ relative to the 2-cell structure H'.

In some future work we plan to investigate the notion of equivalent 2-cell structures, saying that (4.1) is an equivalence whenever (4.2) is, and relate it with homotopy theory.

The notion of a pseudocategory ([**6**],[**5**]) rests in the construction of the induced 2-cells between pullback objects, thus the following definition.

4.2. Cartesian 2-cell structure. It will be useful to consider 2-cell structures such that the functor $H\,(D,\_):\mathbf{C}\longrightarrow Set$ preserves pullbacks for every object D in $\mathbf{C}$, that is: the functor

$$H:\mathbf{C}^{op}\times\mathbf{C}\longrightarrow Set,$$

giving a 2-cell structure to a category $\mathbf{C}$, has the following property

$$H\left(D,A\times_{\{f,g\}}B\right)\overset{\varphi}{\cong}\{(x,y)\in H\,(D,A)\times H\,(D,B)\mid fx=gy\}$$

for every object D in $\mathbf{C}$ and pullback diagram

$$\begin{tikzcd} A\times_C B \arrow[r, "\pi_2"] \arrow[d, "\pi_1"'] & B \arrow[d, "f"] \\ A \arrow[r, "g"] & C \end{tikzcd} ,$$

where φ is required to be a natural isomorphism, that is, for every $h : D \longrightarrow D'$, the following square commutes

$$\begin{tikzcd}
H(D, A \times_C B) \arrow[r, "\cong\varphi"] \arrow[d, "H(h,1)"'] & \{(x, y) \mid fx = gy\} \arrow[d] \\
H(D', A \times_C B) \arrow[r, "\cong\varphi"] & \{(x', y') \mid fx' = gy'\}
\end{tikzcd}$$

or in other words, that

$$\langle x, y\rangle \, h = \langle xh, yh\rangle$$

as displayed in the diagram below

$$\begin{tikzcd}
D' \arrow[r, "h"] & D \arrow[dr, Rightarrow, dashed, "<x,y>"] \arrow[drr, Rightarrow, bend left, "y"] \arrow[ddr, Rightarrow, bend right, "x"'] & & & . \\
& & A \times_C B \arrow[r] \arrow[d] & B \arrow[d, "g"] & \\
& & A \arrow[r, "f"'] & C &
\end{tikzcd}$$

In particular, for $D = A' \times_{C'} B'$, and appropriate x, y, z as in

$$\begin{tikzcd}
A' \arrow[d, Rightarrow, "x"'] & A' \times'_C B' \arrow[l, "\pi'_1"'] \arrow[r, "\pi'_2"] \arrow[d, Rightarrow, "x \times_z y"'] & B' \arrow[d, Rightarrow, "y"] & , & C' \arrow[d, Rightarrow, "z"] & , \\
A & A \times_C B \arrow[l, "\pi_1"'] \arrow[r, "\pi_2"] & B & , & C &
\end{tikzcd}$$

it follows that $x \times_z y$ is the unique element (2-cell) in $H\left(A' \times_{C'} B', A \times_C B\right)$ satisfying

$$\begin{aligned} \pi_2\left(x \times_z y\right) &= y\pi'_2 \\ \pi_1\left(x \times_z y\right) &= y\pi'_1. \end{aligned}$$

Let $\mathbf{C}$ be a category.

Definition 5 (cartesian 2-cell structure). *A 2-cell structure* $(H, \mathrm{dom}, \mathrm{cod}, 0, +)$ *over the category* $\mathbf{C}$ *is said to be* Cartesian *if the functor* $H\left(D, \_\right) : \mathbf{C} \longrightarrow$ *Set preserves pullbacks for every object* D *in* $\mathbf{C}$.

5. Pseudocategories

The notion of pseudocategory (as introduced in [**6**]) is only defined internally to a 2-category. Here we extend it to the more general context of a category with a 2-cell structure (or sesquicategory).

First consider three leading examples.

In any category $\mathbf{C}$, it is always possible to consider two different 2-cell structures, namely the discrete one, obtained when $H = \mathrm{hom}$ and $\mathrm{dom}, \mathrm{cod}, 0, +$ are all identities, and the codiscrete one, obtained when $H = \mathrm{hom} \times \mathrm{hom}$, dom is second projection, cod is first projection, 0 is diagonal and $+$ is uniquely determined. A pseudocategory in the first case is an internal category in $\mathbf{C}$, while in the second case is simply a precategory in $\mathbf{C}$.

In the case of $\mathbf{C} = Cat$, and choosing the natural transformations to be the 2-cell structure, a pseudocategory becomes a pseudo-double-category (see [**6**] and references there), which is at the same time a generalization of a double-category and a bicategory.

At this level of generality, it becomes clear that there is no particular reason why to prefer a specific 2-cell structure in a category instead of another.

For instance, in Top it is usually considered the 2-cell structure obtained from the homotopy classes of homotopies, but others may be consider as well.

Let $\mathbf{C}$ be a category with pullbacks and a cartesian 2-cell structure $(H, \mathrm{dom}, \mathrm{cod}, 0, +)$ defined over it.

Definition 6. *A pseudocategory in* $\mathbf{C}$*, with respect to the 2-cell structure* $(H, \mathrm{dom}, \mathrm{cod}, 0, +)$*, is a system*

$$(C_0, C_1, d, c, e, m, \alpha, \lambda, \rho)$$

where C_0, C_1 *are objects,* d, c, e, m *are morphisms, displayed as*

$$\begin{tikzcd}
C_2 \arrow[r, "m"] & C_1 \arrow[r, shift left=2, "d"] \arrow[r, shift right=2, "d"'] & C_0 \arrow[l, "e" description] & {,\ de = 1 = ce, dm = d\pi_2, cm = c\pi_1}
\end{tikzcd}$$

and α, λ, ρ *are natural and invertible 2-cells, in the sense that*

$$\alpha \in H(C_3, C_1) \text{ and } \lambda, \rho \in H(C_1, C_1)$$

with

$$\begin{aligned}
\mathrm{dom}(\alpha) &= mm_1\ ,\ \mathrm{cod}(\alpha) = mm_2\\
\mathrm{dom}(\lambda) &= me_2\ ,\ \mathrm{dom}(\rho) = me_1\ ,\ \mathrm{cod}(\lambda) = 1_{C_1} = \mathrm{cod}(\rho)
\end{aligned}$$

satisfying the following conditions

$$\begin{aligned}
d\lambda &= 0_d = d\rho\\
c\lambda &= 0_c = c\rho\\
d\alpha &= 0_{d\pi_2 p_2}\ ,\ c\alpha = 0_{c\pi_1 p_1}
\end{aligned}$$

$$\lambda e = \rho e$$

$$m(\alpha \times 0_1) + \alpha(1 \times m \times 1) + m(0_1 \times \alpha) = \alpha(m \times 1 \times 1) + \alpha(1 \times 1 \times m) \tag{5.1}$$

$$m(\rho \times 0_1) + \alpha i_0 = m(0_1 \times \lambda)\,, \tag{5.2}$$

with $C_2, C_3, \pi_1, \pi_2, p_1, p_2$ *obtained by the pullback squares*

$$\begin{tikzcd}
C_2 \arrow[r, "\pi_2"] \arrow[d, "\pi_1"'] & C_1 \arrow[d, "c"] & & C_3 \arrow[r, "p_2"] \arrow[d, "p_1"'] & C_2 \arrow[d, "\pi_1"]\\
C_1 \arrow[r, "d"] & C_0 & & C_2 \arrow[r, "\pi_2"] & C_1
\end{tikzcd}$$

and e_1, e_2, m_1, m_2, i_0 *the following induced morphisms*

$$\begin{aligned} e_1 &= \langle 1, ed\rangle : C_1 \longrightarrow C_2 \\ e_2 &= \langle ec, 1\rangle : C_1 \longrightarrow C_2 \\ m_1 &= 1 \times m : C_3 \longrightarrow C_2 \\ m_2 &= m \times 1 : C_3 \longrightarrow C_2 \\ i_0 &= \langle e_1, e_2\rangle : C_2 \longrightarrow C_3. \end{aligned}$$

Some remarks:

A 2-cell $x \in H(A,B)$ is invertible when there is a (necessarily unique) element

$$-x \in H(A,B)$$

such that $\operatorname{dom}(x) = \operatorname{cod}(-x)$, $\operatorname{cod}(x) = \operatorname{dom}(-x)$ and

$$x + (-x) = 0_{\operatorname{cod}(x)} \ , \ (-x) + x = 0_{\operatorname{dom}(x)};$$

A 2-cell $x \in H(A,B)$ is natural when

$$\operatorname{cod}(x)\, y + x \operatorname{dom}(y) = x \operatorname{cod}(y) + \operatorname{dom}(x)\, y \tag{5.3}$$

for every element $y \in H(X,A)$ for every object X in $\mathbf{C}$.

The 2-cells α, λ, ρ may also be presented as

$$\begin{tikzcd} C_3 \arrow[r, bend left=50, "mm_1"{name=U1}] \arrow[r, bend right=50, "mm_2"'{name=D1}] & C_1 & , & C_1 \arrow[r, bend left=50, "me_2"{name=U2}] \arrow[r, bend right=50, "1"'{name=D2}] & C_1 & , & C_1 \arrow[r, bend left=50, "me_1"{name=U3}] \arrow[r, bend right=50, "1"'{name=D3}] & C_1 & . \arrow[Rightarrow, from=U1, to=D1, "\alpha"] \arrow[Rightarrow, from=U2, to=D2, "\lambda"] \arrow[Rightarrow, from=U3, to=D3, "\rho"] \end{tikzcd}$$

Equations (5.1) and (5.2) correspond to the internal versions of the famous MacLane's coherence pentagon and triangle, presented diagrammatically as follows

$$\begin{tikzcd} & \bullet \arrow[rr, "m(0_{C_1}\times_{C_0}\alpha)"] \arrow[dl, "\alpha(1_{C_1}\times_{C_0}1_{C_1}\times_{C_0}m)"'] & & \bullet \arrow[dr, "\alpha(1_{C_1}\times_{C_0}m\times_{C_0}1_{C_1})"] & \\ \bullet \arrow[drr, "\alpha(m\times_{C_0}1_{C_1}\times_{C_0}1_{C_1})"'] & & & & \bullet \arrow[dll, "m(\alpha\times_{C_0}0_{C_1})"] \\ & & \bullet & & \end{tikzcd} \tag{5.4}$$

$$\begin{tikzcd} \bullet \arrow[rr, "\alpha i_0"] \arrow[dr, "m(0_{C_1}\times_{C_0}\lambda)"'] & & \bullet \arrow[dl, "m(\rho\times_{C_0}0_{C_1})"] \\ & \bullet & \end{tikzcd} \tag{5.5}$$

and restated in terms of generalized elements as

(pentagon)
$$\begin{tikzcd}
& f(g(hk)) \arrow[rr, "f\alpha_{g,h,k}"] \arrow[dl, "\alpha_{f,g,hk}"'] & & f((gh)k) \arrow[dr, "\alpha_{f,gh,k}"] & \\
(fg)(hk) \arrow[drr, "\alpha_{fg,h,k}"'] & & & & (f(gh))k \arrow[dll, "\alpha_{f,g,h}k"] \\
& & ((fg)h)k & &
\end{tikzcd}$$

(midle triangle)
$$\begin{tikzcd}
f(1g) \arrow[rr, "\alpha_{f,1,g}"] \arrow[dr, "f\lambda_{g}"'] & & (f1)g \arrow[dl, "\rho_{f}g"] \\
& fg &
\end{tikzcd}$$

where $m\langle f,g\rangle = fg$.

To check that a given 2-cell $x\in H(A,B)$ is natural is, in general, a complicated task: we have to analyze equation (5.3) for every possible y. On the other hand, removing naturality conditions for $\alpha,\lambda,\rho,$ we loose the Coherence Theorem [**4**] and there is no longer guaranteed that for example the following diagrams are commutative

(5.6)
$$\begin{tikzcd}
1(fg) \arrow[rr, "\alpha_{1,f,g}"] \arrow[dr, "\lambda_{fg}"'] & & (1f)g \arrow[dl, "\lambda_{f}g"] \\
& fg &
\end{tikzcd}$$

(5.7)
$$\begin{tikzcd}
f(g1) \arrow[rr, "\alpha_{f,g,1}"] \arrow[dr, "f\rho_{g}"'] & & (fg)1\ . \arrow[dl, "\rho_{fg}"] \\
& fg &
\end{tikzcd}$$

This diagrams, when internalized, correspond, respectively, to the following equations

$$\begin{aligned}
m\left(\lambda\times 0_{C_1}\right)+\alpha i_2 &= \lambda m,\\
\rho m+\alpha i_1 &= m\left(0_{C_1}\times\rho\right)
\end{aligned}$$

and since the 2-cells are invertible, the above set of equations may be presented as

$$\begin{aligned}
\alpha i_2 &= -m\left(\lambda\times 0_{C_1}\right)+\lambda m,\\
\alpha i_1 &= -\rho m+m\left(0_{C_1}\times\rho\right).
\end{aligned}$$

Now, in the context of a weakly Mal'cev sesquicategory, as it is proved in [**9**], the 2-cell α is uniquely determined by αi_2 and αi_1 and hence it only depends on λ,ρ and m.

This suggests to consider a (non natural) version of a pseudocategory in a sesquicategory.

Let $\mathbf{C}$ be a category with pullbacks and a cartesian 2-cell structure $(H,\mathrm{dom},\mathrm{cod},0,+)$ defined over it.

A (non natural) pseudo category internal to **C** and relative to the 2-cell structure $(H, \mathrm{dom}, \mathrm{cod}, 0, +)$ is a system

$$(C_0, C_1, d, c, e, m, \alpha, \lambda, \rho)$$

where C_0, C_1 are objects, d, c, e, m are morphisms as in the definition above, and, α, λ, ρ are invertible 2-cells $\alpha \in H(C_3, C_1)$ and $\lambda, \rho \in H(C_1, C_1)$ satisfying the following conditions (with $C_2, C_3, \pi_1, \pi_2, p_1, p_2, e_1, e_2, m_1, m_2, i_0$ defined as above)

$$\begin{aligned} \mathrm{dom}(\alpha) &= mm_1 \ , \ \mathrm{cod}(\alpha) = mm_2 \\ \mathrm{dom}(\lambda) &= me_2 \ , \ \mathrm{dom}(\rho) = me_1 \ , \ \mathrm{cod}(\lambda) = 1_{C_1} = \mathrm{cod}(\rho) \end{aligned}$$

$$\begin{aligned} d\lambda &= 0_d = d\rho \\ c\lambda &= 0_c = c\rho \\ d\alpha &= 0_{d\pi_2 p_2} \ , \ c\alpha = 0_{c\pi_1 p_1} \end{aligned}$$

$$\lambda e = \rho e$$

$$m(\alpha \times 0_1) + \alpha(1 \times m \times 1) + m(0_1 \times \alpha) = \alpha(m \times 1 \times 1) + \alpha(1 \times 1 \times m) \tag{5.8}$$

$$\alpha i_0 = -m(\rho \times 0_1) + m(0_1 \times \lambda) \tag{5.9}$$

$$\alpha i_2 = -m(\lambda \times 0_{C_1}) + \lambda m \tag{5.10}$$

$$\alpha i_1 = -\rho m + m(0_{C_1} \times \rho), \tag{5.11}$$

$$\lambda \circ \lambda, \lambda \circ \rho, \rho \circ \rho, \rho \circ \lambda. \tag{5.12}$$

Of course that in the case α, λ, ρ are natural 2-cells then the last three conditions are redundant and we obtain Definition 6 above.

6. Conclusion

We conclude this note by giving three results of application.

The first example is an instance of section 3.1, the second example is an application of section 3.2, while the third one is from 3.4.

In the setting of section 3.1, let **B** be a weakly Mal'cev category (see [**8**], or simply assume that **B** is a Mal'cev category) and consider **C** =Cat(**B**) and $(H, \mathrm{dom}, \mathrm{cod}, 0, +)$ as in section 3.1.

In particular, **C** as above with internal transformations (not necessarily natural), is a weakly Mal'cev sesquicategory and the following result is proved in [**9**].

Theorem 1. *A (non natural) pseudo category internal to* **C** *and relative to the specific 2-cell structure as above, satisfying the additional condition*

$$\lambda e = 0_e = \rho e$$

is completely determined by a reflexive graph in **C**

$$\begin{tikzcd} C_1 \arrow[r, shift left=2, "d"] \arrow[r, shift right=2, "c"'] & C_0 \arrow[l, "e" description] \end{tikzcd} \quad , \quad de = 1_{C_0} = ce$$

together with 2-cells

$$\lambda, \rho \in H(C_1, C_1)$$

satisfying the following conditions,

$$\mathrm{cod}(\lambda) = 1_{C_1} = \mathrm{cod}(\rho)$$

$$\begin{aligned} d\lambda &= 0_d = d\rho \\ c\lambda &= 0_c = c\rho \\ \lambda e &= 0_e = \rho e \end{aligned}$$

and furthermore it is equipped with a morphism

$$m : C_2 \longrightarrow C_1$$

uniquely determined by

$$me_1 = u \ , \ me_2 = v$$

and a 2-cell $\alpha \in H\left(C_3, C_1\right)$*, uniquely determined by*

$$\alpha i_1 e_1 = -\rho u \ , \ \alpha i_2 e_1 = -u\lambda + \lambda u \ , \ \alpha i_2 e_2 = \lambda v$$

where $v = \operatorname{dom}\left(\lambda\right)$ *and* $u = \operatorname{dom}\left(\rho\right)$*.*

Note that this result reflects the striking fact that, in this case,

$$5.10 + 5.11 + 5.12 \Rightarrow 5.8 + 5.9,$$

and if α, λ, ρ are natural then the description above is in fact a pseudocategory.

The second example describes pseudocategories in Groups, and it gives the notion of crossed module with the freedom to choose an element δ in the centre of X.

If considering the category of Groups with the 2-cells structure given by derivations as in 3.2 then an internal pseudocategory is completely determined by a group homomorphism

$$X \xrightarrow{\partial} B,$$

an action of B in X (denoted by $b \cdot x$) and a distinguished element in X, δ satisfying the following conditions

$$\begin{aligned} \partial\delta &= 0 \\ x &= \delta + x - \delta \\ \partial\left(b \cdot x\right) &= b\partial\left(x\right) b^{-1} \\ \partial\left(x\right) \cdot \bar{x} &= x + \bar{x} - x. \end{aligned}$$

Where the objects are the elements of B, the arrows are pairs $(x, b) : b \longrightarrow \partial x + b$ and the composition of $(x', \partial x + b) : \partial x + b \longrightarrow \partial x' + \partial x + b$ with $(x, b) : b \longrightarrow \partial x + b$ is $(x' + x - \lambda + b \cdot \lambda, b) : b \longrightarrow \partial x' + \partial x + b$. The isomorphism between $(0, \partial x + b) \circ (x, b) = (x, b) \circ (0, b)$ and (x, b) is the element $(\delta, 0) \in X \rtimes B$. Associativity is satisfied, since $(x'', \partial x' + \partial x + b) \circ ((x', \partial x + b) \circ (x, b)) = ((x'', \partial x' + \partial x + b) \circ (x', \partial x + b)) \circ (x, b)$.

In the case of an additive category $\mathbf{A}$ with kernels and a 2-cell structure given by an Ab-functor

$$H : \mathbf{A}^{op} \times \mathbf{A} \longrightarrow Ab$$

and a natural transformation

$$D : H \longrightarrow \hom_A$$

and using the notation

$$[x, y] = D\left(x\right) y - xD\left(y\right)$$

a pseudo category is completely determined by

$$A \xrightarrow{h} B$$

$$\begin{aligned}\lambda,\rho &\in H(A,A)\\ \eta &\in H(B,A)\end{aligned}$$

$$\begin{aligned}h\lambda &= 0\\ h\rho &= 0\\ h\eta &= 0\end{aligned}$$

where α is given by $(5.9), (5.10)$ and (5.11) and (5.8) translates to

$$\begin{gathered}(1-D\rho)[\rho,\rho]=0\\ D\lambda[\rho,\rho]=D\rho[\lambda,\rho]+(1-D\rho)[\rho,\lambda]\\ (1-D\lambda)[\lambda,\rho]+D\lambda[\rho,\lambda]=D\rho[\lambda,\lambda]+(1-D\rho)[\rho,\eta]h\\ (1-D\lambda)[\lambda,\lambda]=(1-D\lambda)[\lambda,\eta]\\ (1-D\rho-D\lambda)[\lambda,\eta]=(1-D\rho-D\lambda)[\rho,\eta]\end{gathered}$$

which is trivial as soon as we introduce (5.12).

The pseudocategory thus determined is of the form (see [**5**])

$$\begin{tikzcd}
A\oplus A\oplus B \arrow[r, "m"] & A\oplus B \arrow[r, shift left=2, "(0\ 1)"] \arrow[r, shift right=2, "(\delta\ 1)"'] & B \arrow[l, "\binom{0}{1}"']
\end{tikzcd}$$

$$m=\begin{pmatrix} f & g & h\\ 0 & 0 & 1\end{pmatrix}$$

$$\begin{aligned}g &= 1-D(\lambda)\\ f &= 1-D(\rho)\\ h &= -D(\eta)\end{aligned}$$

$$\alpha=\begin{pmatrix}\alpha_1 & \alpha_2 & \alpha_3 & \alpha_0\\ 0 & 0 & 0 & 0\end{pmatrix}$$

$$\begin{aligned}\alpha_1 &= -f\rho\\ \alpha_2 &= \lambda+g\rho-\rho-f\lambda\\ \alpha_3 &= g\lambda-f\eta\delta\\ \alpha_0 &= g\eta-f\eta\end{aligned}$$

$$\lambda=\begin{pmatrix}\lambda & \eta\\ 0 & 0\end{pmatrix}\qquad\qquad \rho=\begin{pmatrix}\rho & \eta\\ 0 & 0\end{pmatrix}.$$

In particular the category of abelian chain complexes is of this form.

Another example of this form is the case of TAG, Topological Abelian Groups.

Let $\mathbf{C}=TAG$ the category of topological abelian groups with the 2-cell structure given by

$$H(X,Y)=\{\alpha: I\times X\longrightarrow Y \mid \alpha \text{ is continuous, } \alpha(0,\_)=0 \text{ and } \alpha(t,\_) \text{ is a homomorphism}\}/\sim$$

where I is the unit interval and the equivalence $\sim$ is defined by

$$\alpha\sim\beta$$

if and only if $\alpha(1,\_) = \beta(1,\_) = h$, there is $\Phi : I \times I \times X \longrightarrow Y$, continuous and such that

$$\begin{aligned} \Phi(0,\_,\_) &= \alpha \\ \Phi(1,\_,\_) &= \beta \\ \Phi(s,0,\_) &= 0 \\ \Phi(s,1,\_) &= h \end{aligned}$$

and $\Phi(s,t,\_)$ is a homomorphism.

The natural transformation $D : H \longrightarrow \hom$ is given by

$$D([\alpha]) = \alpha(1,\_)$$

with

$$(g[\alpha]f)(t,x) = g(\alpha(t,f(x))).$$

We have

$$[\alpha]D([\beta]) = D([\alpha])[\beta]$$

because

$$\alpha D(\beta) \sim D(\alpha)\beta \Leftrightarrow \alpha(t,\beta(1,x)) \sim \alpha(1,\beta(t,x))$$

and there is

$$\Phi(s,t,x) = \alpha\left(t^{(1-s)}, \beta(t^s,x)\right).$$

From [**5**] we now conclude that a pseudocategory in TAG (with the 2-cell structure as above) is given by a morphism in TAG

$$k : A \longrightarrow B,$$

together with

$$\lambda, \rho : I \times A \longrightarrow A$$

in $H(A,A)$ and also

$$\eta : I \times B \longrightarrow A$$

in $H(B,A)$ satisfying

$$k\rho(t,\_) = 0, k\lambda(t,\_) = 0, k\eta(t,\_) = 0.$$

The objects in the pseudocategory are the points in B while the pseudomorphisms are pairs (a,b) with domain b and codomain $k(a)+b$; the composition of

$$b \xrightarrow{(a,b)} b' \xrightarrow{(a',b')} b''$$

is given by the following formula

$$(a - \rho(1,a) + a' - \lambda(1,a') - \eta(1,b), b).$$

In particular, if we consider that A is the space of paths in B starting at zero

$$A = \{x : I \longrightarrow B \mid x \text{ continuous and } x(0) = 0\}$$

with

$$k(x) = x(1)$$

and choose representatives of λ, ρ and η as

$$\begin{aligned}
\lambda\left(s,x\right)\left(t\right) &= \begin{cases} x\left(st\right)-x\left(2st\right) & \text{if} \quad t\leqslant\frac{1}{2} \\ x\left(st\right)-x\left(s\right) & \text{if} \quad t\geqslant\frac{1}{2} \end{cases} \\
\rho\left(s,x\right)\left(t\right) &= \begin{cases} x\left(st\right) & \text{if} \quad t\leqslant\frac{1}{2} \\ x\left(st\right)-x\left(2st-s\right) & \text{if} \quad t\geqslant\frac{1}{2} \end{cases} \\
\eta &= 0
\end{aligned}$$

then we obtain the usual composition of paths

$$y+x=\begin{cases} x\left(2t\right) & \text{if} \quad t\leqslant\frac{1}{2} \\ y\left(2t-1\right)+x\left(1\right) & \text{if} \quad t\geqslant\frac{1}{2} \end{cases} .$$

References

[1] Crans, S.E: A tensor product for Gray-categories, Theory and Applications of Categories, Vol.5, N^{o}.2, 1999, pp.12-69.

[2] Gray, J.W.: *Formal Category Theory: Adjointness for 2-categories*, Lecture Notes in Mathematics, Springer-Verlag, 1974

[3] Leinster, T.: *Higher Operads, Higher Categories*, London Mathematical Society Lecture Notes Series, Cambridge University Press, 2003 (electronic version).

[4] MacLane, S.: *Categories for the working Mathematician*, Springer-Verlag, 1998, 2^{nd} edition.

[5] Martins-Ferreira, N.: Internal Weak Categories in Additive 2-Categories with Kernels, Fields Institute Communications, Volume 43, p.387-410, 2004.

[6] Martins-Ferreira, N: Pseudo-categories, *JHRS*, Vol.1(1), 2006, pp.47-78.

[7] Martins-Ferreira, N.: The (tetra)category of pseudocategories in additive 2-categories with kernels, Applied Categorical Structures (published in Online First, and to appear in print), 2008.

[8] Martins-Ferreira, N.: Weakly Mal'cev categories, Theory and Applications of Categories, Vol. 21, No. 6, pp 91-117, 2008.

[9] Martins-Ferreira, N.: Low-dimensional internal categorical structures in weakly Mal'cev sesquicategories, PhD Thesis, 2008.

[10] Street, R.H.: Cosmoi of internal categories, Trans. Amer. Math. Soc. 258, 1980, 271-318

[11] Street, R.H.: Fibrations in Bicategories, Cahiers Topologie et Géométrie Diferéntielle Catégoriques, 21:111-120, 1980.

[12] Street, R.H.: *Handbook of Algebra*, chapter Categorical Structures, pages 529–577. Elsevier Science, 1996.

[13] Bourn, D., "Another denormalization theorem for the abelian chain complexes", J. Pure and Applied Algebra 66, 1990, 229-249.

[14] Brown, R. and Higgins, Ph.J., "Cubical Abelian Groups with Connections are Equivalent to Chain Complexes", Homology, Homotopy and Applications vol. 5(1), 2003, 49-52.

Polytechnic Institute of Leiria
E-mail address: `nelsonmf@estg.ipleiria.pt`
URL: `http://www.estg.ipleiria.pt/~nelsonmf`